



The influence of macronutrient intake, stress and prostaglandin levels (pgf2 α) of urine with the incidence of dysmenorrhea in adolescents

Asriani Tahir^a, Andi Wardihan Sinrang^{a,*}, Elizabeth Catherine Jusuf^b, Saidah Syamsuddin^c, Stang^d, Aryadi Arsyad^a

^a Midwifery Department, Graduate School, Universitas Hasanuddin, Indonesia

^b Department of Obstetrics and Gynaecology, Faculty of Medicine, Universitas Hasanuddin, Indonesia

^c Department of Psychiatry, Faculty of Medicine, Universitas Hasanuddin, Indonesia

^d Department of Biostatistics, Faculty of Public Health, Universitas Hasanuddin, Indonesia

ARTICLE INFO

Article history:

Received 28 June 2021

Accepted 30 July 2021

Keywords:

Intake of macronutrients

Stress

Prostaglandin levels (pgf2 α)

Dysmenorrhea

ABSTRACT

Objectives: This study aimed to analyze the influence of macronutrient intake, stress, and prostaglandin levels (pgf2 α) on adolescent dysmenorrhea incidence.

Method: This type of study is observational analytic with a cohort study draft done in January–March 2020 at High junior school 21 Makassar. Respondents in this study were grade X and XI students divided into 64 teenagers who had dysmenorrhea and 64 adolescents who did not experience Dysmenorrhea. The criteria of the respondent in this study were the reproductive age, already experiencing menstruation, knowing the time and date of menstruation, menstrual cycles were regular, and willing to be respondents. The study used *Menstrual Symptoms Questionnaire (MSQ)* and used an ultrasonography (ultrasound) examination to perform the sample cervical. *Food recall* 24 hours to assess the intake of macronutrients, *Depression Anxiety Stress Scales (DASS 42)* to measure stress levels, and an examination of urine prostaglandin levels using the method Enzyme-Linked Immunosorbent Assay (ELISA). Urine intake is carried out on the second day as much as 2–5 cc. Data were analyzed by the Chi-square test and logistics regression backward.

Result: A multivariate analysis showed a variable that strongly affects dysmenorrhea is stress with the value $p = 0.000$ and the level of prostaglandins with $p\text{-value} = 0.003$ compared to other variables.

Conclusion: Stress and prostaglandin levels significantly affect the occurrence of dysmenorrhea in adolescents.

© 2021 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

During menstruation, women experience a variety of disorders that can cause physical and disturbing activities. One inconvenience is dysmenorrhea. Dysmenorrhea is a public health problem that means menstruation that nyeri.^{1–4}

Dysmenorrhea is a health problem that has a negative impact on physical and emotional health and most causes absence in school that affects academic presentation. Dysmenorrhea in women can affect students' academic abilities, school absences, and loss of daily work.^{5–7} A dysmenorrhea incidence rate in Indonesia amounted to 107,673 people, consisting of 59,671 people experiencing primary dysmenorrhea (54.89%) The soul and 9496 people experience secondary dysmenorrhea (9.36%) Jiwa.^{8,9}

Balanced dietary nutritional intake (e.g., due to protein or high fat, or low fiber intake) is associated with a greater risk of obesity in children and adolescents than adults. Therefore, it is not only the macro intake needed for health, but the quality of the diet also affects examples of fats and carbohydrates (complex and straightforward fibers and carbohydrates) and proteins (animals

and plants). Nutritional Status of overweight and obesity is also a risk factor of dysmenorrhea.^{10,11}

The pain that accompanies uterine contractions can affect functional mechanisms, and one of the causes of dysmenorrhea is psychic, and one of those psychic factors is the stress that causes a physiological stress response. The stress response includes activating the sympathetic nervous system and the release of various hormones and peptides. More and more formed prostaglandins and vasopressin cause the uterine contraction of the uterus to clamp the ends of the nerve fibers; further, the formation is transmitted through sympathetic nerve fibers.^{12–14}

Methods

Research sites

The study was conducted in January–March 2020 and has received a recommendation of ethical approval with the protocol number UH19110996. Research conducted at High junior school 21 Makassar and laboratory of Microbiology of the Hasanuddin University of Makassar.

Data types and sources

The Data collected from the samples are age, menstrual cycle, prolonged menstruation, macronutrient intake, stress level, urine prostaglandin levels, and medical diagnosis. Data were taken from the students who are dysmenorrhea and not dysmenorrhea.

Peer-review under responsibility of the scientific committee of the 3rd International Nursing, Health Science Students & Health Care Professionals Conference. Full-text and the content of it is under responsibility of authors of the article.

* Corresponding author.

E-mail addresses: wardihan@pasca.unhas.ac.id, pmc@agri.unhas.ac.id (A.W. Sinrang).

Table 1

Characteristics of the respondents group of cases and controls on high junior school 21 Makassar.

Characteristics	n = 128	%
Age		
14 years	1	0.8
15 years	35	27.3
16 years	74	57.8
17 years	18	14.1
Menstrual cycle		
<21 week	3	2.3
21–35 week	124	96.9
>35 week	1	0.8
Menstruation		
<5 day	19	14.9
5–7 day	89	69.5
7–14 day	20	15.6

Source: Primary Data year 2020.

Data collection technique

Data related to respondents were gathered through questionnaires, live interviews, and the respondent's USG examination and urine sampling. The collected samples are then stored in the refrigerator at a temperature of -20°C . After the samples were fulfilled, the examination of prostaglandin (pgf2 α) used the Kit Elisa, Bioassay Technology Laboratory at the Hasanuddin University of Makassar Hospital laboratory.

Results

The majority of respondents were 16 years old with menstrual cycles of 21–35 weeks and menstrual periods of 5–7 days (Table 1).

Most respondents have less intake of macronutrients (carbohydrates, protein, fat). Most respondents have moderate stress levels, and respondents have the lowest levels of prostaglandins. The number of respondents who experienced dysmenorrhea was 64 people and not dysmenorrhea 64 people (Table 2).

Respondents who experience dysmenorrhea and not dysmenorrhea have less macronutrient intake (carbohydrate, fat, protein), i.e., the p -value is >0.05 (Table 3).

Respondents who experienced dysmenorrhea had the most moderate stress, namely, 30 respondents (46.9%), and respondents who did not experience dysmenorrhea had the most normal stress, namely 26 respondents (0.0%). Chi-square test results obtained $p=0.002$ ($p < 0.05$). This shows there is a relationship between stress with the incidence of dysmenorrhea (Table 4).

Prostaglandin levels with p -value 0.002, which means <0.005 . This shows that the incidence of dysmenorrhea influences prostaglandin (pgf2 α) levels (Table 5).

Table 3

Influence of macronutrient intake with dysmenorrhea incidence.

Variable	Dysmenorrhea (n %)		p-value	RR value	CI 95%	
					Lower	Upper
Carbohydrate	Less	38 (59.4%)	0.462 ^a	1.259	0.805	1.968
	More	12 (18.8%)		1.016	0.571	1.808
	Good	14 (21.9%)		1.000		
Protein	Less	39 (60.9%)	0.093 ^a	1.159	0.796	1.687
	More	5 (7.8%)		0.569	0.254	1.278
		20 (31.3%)		1.000		
Fat	Less	36 (56.3%)	0.0615	1.092	0.695	1.716
	More	14 (21.9%)		1.292	0.772	2.161
	Good	14 (21.9%)		1.000		

^a Chi-square.

Table 2

Variable case group and control study on high junior school 21 Makassar.

Variable	n = 128	%
Macronutrient Intake		
<i>a. Carbohydrates</i>		
Less	69	53.9
More	27	21.1
Good	32	25.0
<i>b. Protein</i>		
Less	69	53.9
More	18	14.1
Good	41	32.0
<i>c. Fat</i>		
Less	73	57.0
More	24	18.8
Good	31	24.2
Stress		
Normal	36	28.1
Mild	36	28.1
Are	42	32.8
Severe	13	10.2
Very severe	1	0.8
Prostaglandins		
Low (<181.9 pg/ml)	85	66.4
High (≥ 181.9 pg/ml)	43	33.6
Dysmenorrhea		
Yes	64	50
Not	64	50

Source: Primary Data year 2020.

Variables that are very influential on dysmenorrhea are stress with a value of $p = 0.000$ and prostaglandins with a value of $p = 0.003$ compared to other variables) (Table 6).

Discussion

Table 1 shows that most 16-year-old are as many as 74 people (57.8%). Respondents with the most menstrual cycles are 21–35 days of 124 people (96.9%). Respondents with the most menstrual period of 5–7 days are 89 people (69.5%). Table 2 shows that respondents who have the highest intake of carbohydrate macronutrients are less than 69 people (53.9%), the maximum protein intake of macronutrients is less than 69 people (53.9%), consumption of the fattest macronutrient intake is less than 73 people (57.0%). Most of the Respondents have low prostaglandin levels of 66.4%. Previously, the cut-off point was determined at prostaglandin levels (pgf2 α) using a ROC curve with 50.00% sensitivity and 82.81% specificity values. Respondents who have been dysmenorrhea as much as 64 (50%) and did not suffer from dysmenorrhea as much as 64 respondents (50%).

Carbohydrates are one part of the macronutrients contained in foods. High protein levels in foods will increase the levels of tyrosine that implicate the increase in the synthesis of dopamine, the

Table 4

The impact of stress analysis with dysmenore events.

Variable		Dysmenorrhea (n %)	Not dysmenorrhea (n %)	p-value	RR value	CI 95%	
						Lower	Upper
Stress	Low	15 (23.4%)	21 (32.8%)	0.002 ^a	1.5	0.78	2.883
	Medium	30 (46.9%)	12 (18.8%)		2.571	1.468	4.503
	High	8 (12.5%)	5 (7.8%)		2.215	1.123	4.372
	Awfully	1 (1.6%)	0 (0.0%)		3.6	2.126	6.096
	Normal	10	26		1		

^a Chi-square.

Table 5

Influence of prostaglandins (pgf2α) levels with dysmenorrhea events.

Variable	B	S.E.	Wald	df	p-value	OR	95% CI for OR	
							Lower	Upper
Prostaglandin	0.030	0.010	9.642	1	0.002 ^a	1.031	1.011	1.050

^a Regression logistic.

Table 6

Influence analysis of macronutrient intake, stress, prostaglandin levels (Pgf2α) with dysmenorrhea incidence.

	B	S.E.	Wald	df	p-value	OR	95% CI for OR	
							Lower	Upper
Step 1 ^a	Stress	1.489	0.4	13.818	0.000	4.431	2.021	9.713
	Prostaglandin	0.031	0.01				1.032	1.011

^a Regression logistic backward.

dopamine metabolite. Dopamine is purported to improve satisfaction and mood. In general, a recognized breakfast contributes positively to the quality of the diet, the intake of macronutrients and micronutrients, body mass index, lifestyle, and positively affects behavioral, cognitive, and academic performance in children aged-school.^{15,16} The results of this study with a Chi-square test showed no influence between the intake of macronutrients (carbohydrates, proteins, fats) with the incidence of dysmenorrhea.

Fatty acids are an essential component in cell membranes as they can produce energy. Therefore, women should be consuming foods high in omega-3 fatty acids, e.g., fish (salmon, tuna, mackerel, dried fish), fish oil, soy, eggs, meat, shrimp, and fruits. Conversely, consuming foods that are low in omega-3 can increase pain dysmenorrhea.¹⁷ The results of this study differ from those done in Georgia, stating that energy and fat intake are connected with dysmenorrhea events.

Tables 3 and 4 indicates that there is a stress effect with the occurrence of Dysmenore with the value $p=0.002$. Stress can exacerbate the current state of menstruation. The research is in line with India's research that there is a significant relationship between dysmenorrhea and stress. In addition, the Korean nursing Program recommends that teenagers improve their health and encourage them to eat regularly to reduce stress when menstruation.¹⁸

The cause of pain is dysmenorrhoea due to an increase in prostaglandin hormones. This hormone results in uterine contractions and vascular vasoconstriction. Blood flow that leads to the uterus decreases so that the uterus does not get adequate oxygen, causing pain.¹⁹ In line with the test results in the 3.5 table indicates that the incidence of dysmenorrhea has an influence on the level of prostaglandins (Pgf2α) with a value $p=0.002$ (<0.005).

A multivariate analysis using a backward logistics regression equation found that a variable that strongly affects dysmenorrhea is stress with the value $p=0.000$ and prostaglandins with a value $p=0.003$ compared to other variables. Increased levels of prostaglandins proved to have been found in menstrual fluid in women with severe dysmenorrhea, especially during the first two days of menstruation. Painful menstruation affects the biochemical

and cellular processes throughout the body, including the brain and psychological. During stress, the body will produce excessive adrenal, estrogen, progesterone, and prostaglandin hormones. Increased estrogen hormones may increase excessive uterine contractions. In addition, increased adrenal hormones can cause muscle tension of the uterus, which makes excessive contraction so which will cause a taste of pain.^{20,21}

Conclusion

Levels of prostaglandins (pgf2α) affect the incidence of dysmenorrhea in the event of a change in the period of menstrual hormones, i.e., prostaglandins can reduce or temporarily inhibit the blood supply to the uterus, causing the uterus to lack oxygen and cause myometrium contraction so that there is pain. Furthermore, stress is very influential with dysmenorrhea, among other variables, because of the more stressful one in the face of a problem than the pain that is felt when menstruation is increasing so that stress can interfere with the work of the endocrine system. A person with an insufficient intake of nutrients has a higher risk of being affected by Dysmenore. For that, teenagers must meet the intake of macronutrients consisting of carbohydrates, proteins, and fats, especially during menstruation.

Conflicts of interest

The authors declare no conflict of interest.

References

1. Acheampong K, Baffour-Awuah D, Ganu D. Prevalence and predictors of dysmenorrhea, its effect, and coping mechanisms among adolescents in Shai Osudoku District, Ghana. Obstet Gynecol Int. 2019;2019.
2. Al-Matouq S, Al-Mutairi H, Al-Mutairi O. Dysmenorrhea among high-school students and its associated factors in Kuwait. BMC Pediatr. 2019;19:1–12.
3. Chen CX, Draucker CB, Carpenter JS. China supply custom plastic crate storage crates products cheap price-Zhongyi Plastic Technology Co., Ltd. BMC Womens Health. 2018;18.

4. Harris C, Flexeder C, Thiering E. Changes in dietary intake during puberty and their determinants: results from the GINIplus birth cohort study. *BMC Public Health.* 2015;15:1–19.
5. Aboushady RM-N, El-Saidy TMK. Effect of home based stretching exercises and menstrual care on primary dysmenorrhea and premenstrual symptoms among adolescent girls. *IOSR J Nurs Heal Sci.* 2016;5:10–7.
6. Hailemeskel S, Demissie A, Assefa N. Primary dysmenorrhea magnitude, associated risk factors, and its effect on academic performance: evidence from female university students in Ethiopia. *Int J Womens Health.* 2016;8:489.
7. Shehata NAA, Arafa AE, El Wahed HAA, Fahim AS, Hussein GK. Epidemiology of dysmenorrhea among university students in Egypt. *Int J Womens Heal Wellness.* 2018;4:73.
8. Orimadegun BE, Awolude OA, Agbedana EO. Markers of lipid and protein peroxidation among Nigerian university students with dysmenorrhea. *Niger J Clin Pract.* 2019;22:174.
9. Ramli N, Santy P. Efektifitas Pemberian Ramuan Jahe (*Zingibers officinale*) dan Teh Rosella (*Hibiscus sabdariffa*) terhadap Perubahan Intensitas Nyeri Haid. *AcTion Aceh Nutr J.* 2017;2:61–6.
10. Khodakarami B, Masoomi SZ, Faradmal J. The severity of dysmenorrhea and its relationship with body mass index among female adolescents in Hamadan, Iran. *J Midwifery Reprod Heal.* 2015;3:444–50.
11. López-Sobaler AM, Aparicio A, Rubio J. Adequacy of usual macronutrient intake and macronutrient distribution in children and adolescents in Spain: a National Dietary Survey on the Child and Adolescent Population, ENALIA 2013–2014. *Eur J Nutr.* 2019;58:705–19.
12. Rejeki S, Eldaniati E. Intensitas Nyeri Haid Primer Melalui Senam Dismenore Pada Mahasiswa Di Pondok Pesantren K.H. Sahla Rosjidi Universitas Muhammadiyah Semarang. In: Prosiding Seminar Nasional Unimus. 2018. p. 183–93.
13. Sumiash S, Ningrum P. Efektifitas Pemberian Vitamin E Terhadap Penurunan Dismenore pada Siswi Di SMA Negeri 1 Gading Rejo. *Wellness Heal Mag.* 2019;1:15–21.
14. Salim HHS, Yuniyanti B. Hubungan Tingkat Stres Dengan Tingkat Dysmenorrhoea Pada Siswi Kelas X Dan XI SMK Bhakti Karyakota Magelang Tahun 2014. *J Kebidanan.* 2015;4:51–6.
15. Almatsier S. Prinsip dasar ilmu gizi. Jakarta: Gramedia Pustaka Utama; 2010.
16. Cantika HAK, Ardiaria M, Syauqy A. Peran makronutrien dalam sarapan terhadap performa kognitif remaja. *J Kedokt Diponegoro.* 2017;6:611–21.
17. Famimah F, Margawati A, Fitrianti DY. Hubungan Konsumsi Asam Lemak Omega-3, Aktivitas Fisik Dan Persen Lemak Tubuh Dengan Tingkat Dismenore Pada Remaja. *J Nutr Coll.* 2017;6:268–76.
18. Jeon GE, Cha NH, Sok SR. Factors influencing the dysmenorrhea among Korean adolescents in middle school. *J Phys Ther Sci.* 2014;26:1337–43.
19. Novadela NIT, Hardini RA, Mugiatni M. Perbandingan terapi air putih dengan kompres hangat terhadap penurunan skala nyeri haid (dismenore primer) pada remaja. *J Ilm Keperawatan Sai Betik.* 2019;14:219–25.
20. Mou L, Lei W, Chen J, Zhang R, Liu K, Liang X, et al. Mediating effect of interpersonal relations on negative emotions and dysmenorrhea in female adolescents. *Gen Psychiatry.* 2019;32.
21. Unsal A, Ayrancı U, Tozun M, Arslan G, Calik E. Prevalence of dysmenorrhea and its effect on quality of life among a group of female university students. *Ups J Med Sci.* 2010;115:138–45.