

Original article

Impact of the COVID-19 pandemic on the psychological well-being of adolescents by gender identity

Cinta Folch^{a,b,c,*}, Fabiana Ganem^{a,b,d}, Andreu Colom-Cadena^{a,b}, Isabel Martínez^{a,b}, Carmen Cabezas^e, Jordi Casabona^{a,b,c,d}^a Centre for Epidemiological Studies of Sexually Transmitted Disease and AIDS in Catalonia (CEEISCAT), Badalona, Barcelona, Spain^b Germans Trias i Pujol Research Institute (IGTP), Badalona, Barcelona, Spain^c CIBER de Epidemiología y Salud Pública (CIBERESP), Spain^d Departament de Pediatria, d'Obstetrícia i Ginecologia i de Medicina Preventiva i de Salut Pública, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, Spain^e Public Health Secretary, Health Department of the Government of Catalonia, Barcelona, Spain

ARTICLE INFO

Article history:

Received 5 May 2023

Accepted 26 October 2023

Available online 25 November 2023

Keywords:

COVID-19

Psychological well-being

Adolescents

Gender identity

Schools

Cross-sectional studies

ABSTRACT

Objective: To describe the impact of the COVID-19 pandemic on the psychological well-being of adolescents in Catalan schools by gender identity, and to compare coping strategies adopted to manage the health crisis and their relationship with the self-perceived impact of COVID-19 on mental health.

Method: Cross-sectional study in educational centres that includes 1171 adolescents over 15 years old from October to November 2021. Multivariate logistic regression models were built to evaluate the association between coping strategies with self-perceived impact of the pandemic on mental health.

Results: A greater proportion of girls perceived a worsening in mental health than boys due to COVID-19 (36.9% and 17.8%, respectively). The main emotions reported for both girls and boys were worry and boredom. The study found an association between positive coping strategies with less adverse mental health among girls, whereas unhealthy habits were associated with a higher probability of declaring worsening of mental health for both girls and boys.

Conclusions: This study demonstrated the negative impact of the COVID-19 pandemic on psychological well-being in adolescents and a clearly worse impact on girls. It is important to keep monitoring the medium- and long-term secondary impacts of the pandemic on mental health outcomes of adolescents and to gather information that can improve services for the development of healthy coping strategies during health crises like COVID-19, which include gender perspective.

© 2023 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Impacto de la COVID-19 en el bienestar emocional de adolescentes según identidad de género

RESUMEN

Objetivo: Describir el impacto de la COVID-19 en el bienestar emocional de adolescentes en Cataluña y comparar las estrategias de afrontamiento adoptadas durante la pandemia y su relación con el impacto autopercebido de la COVID-19 en la salud mental según identidad de género.

Método: Estudio transversal en centros educativos de Cataluña incluyendo 1171 alumnos/as mayores de 15 años, de octubre a diciembre de 2021. Mediante modelos de regresión logística multivariante se evaluó la asociación entre las estrategias de afrontamiento y el impacto autopercebido de la pandemia en la salud mental.

Resultados: Una mayor proporción de chicas que de chicos percibía un empeoramiento en su salud mental por la COVID-19 (36,9% y 17,8%, respectivamente). Las principales emociones reportadas tanto en chicas como en chicos fueron preocupación y aburrimiento. Se observó una asociación entre estrategias de afrontamiento positivas y una menor probabilidad de declarar un empeoramiento en la salud mental a raíz de la pandemia en las chicas, mientras que otras estrategias relacionadas con hábitos poco saludables se asociaron a una mayor probabilidad de declarar un empeoramiento de la salud mental en chicas y chicos.

Conclusiones: Este estudio demuestra el impacto negativo de la pandemia de COVID-19 en el bienestar psicológico de los/las adolescentes, especialmente en las chicas. Es importante seguir monitorizando el impacto de la COVID-19 a medio y largo plazo en la salud mental de los/las adolescentes y disponer de información que pueda mejorar los servicios para el desarrollo de estrategias de afrontamiento saludables durante crisis de salud como la COVID-19 considerando una perspectiva de género.

© 2023 SESPAS. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Palabras clave:

COVID-19

Bienestar psicológico

Adolescentes

Identidad de género

Escuelas

Estudio transversal

* Corresponding author.

E-mail address: cfolch@iconcologia.net (C. Folch).

Introduction

The SARS-CoV-2 emergency affected health systems globally, compelling governments to implement strategies, like social distancing or confinements, and generating stress, anxiety, and depression which impacted the population well-being, and young people especially.¹ In educational settings, measures such as suspending classes and accelerating the use of technology for learning, had an impact on the routines of students and their families.² Additionally, social distancing reduced the interpersonal interactions and leisure activities of adolescents, who lost social relations with peers, amongst other changes.³ Previous systematic reviews have demonstrated that mental health problems in children and adolescents increased during the pandemic compared to pre-pandemic data, being gender, socio-economic status, previous state of mental and physical health, social support, and consistent routines among the most reported influencing factors.⁴⁻⁸

Several studies have shown gender differences in the response of adolescents to the COVID-19 pandemic, with a worse effect on psychological well-being observed in girls.^{9,10} Reasons cited for the gender differences include a reduction in emotional support from social networks due to social distancing¹¹ and a greater increase in unhealthy lifestyle habits compared to boys.⁹ Additionally, depression and anxiety increase during puberty, especially among girls,¹² which may partially explain these gender disparities. The pandemic highlighted gender differences in mental health that are a reflection of the distinct social processes for youth with different gender identities, as well as disparities in mental health that emerge and intensify during this stage of life.¹³

Effective coping strategies, defined as cognitive and behavioural efforts made to manage situations that are potentially threatening or stressful,¹⁴ have been reported in adolescents, such as physical activity, use of social networks and establishing routines.^{15,16} However, unhealthy coping strategies such as consumption of tobacco, alcohol and/or drugs, or unhealthy eating habits have been identified, although to a lesser extent.¹⁷ Evidence indicates that positive coping strategies, such as connecting with friends and family members and/or engaging in physical activities, are closely related to better mental health outcomes during the COVID-19 pandemic,⁸ whereas unhealthy coping strategies relate more strongly to greater negative impacts on stress and mental health. Also gender is one of the factors that influences which coping strategies are adopted to confront stress.^{10,18} A previous study in Spain on a sample of university students showed that men resorted to physical exercise during the pandemic, while women listened to more music and watched more television.¹⁹

The COVID-19 Sentinel Schools Network of Catalonia (CSSNC), main objective was to monitor and evaluate the COVID-19 situation and its impact in educational settings in order to gather evidence for the health policies, prevention and control of SARS-CoV-2 in schools. Since its implementation in 2020, mental health was identified as a priority challenge, leading us to include in the project new research questions to better explore this issue in schools and develop more effective strategies.²⁰

This study aims to describe the impact of the COVID-19 pandemic on psychological well-being, defined as “a state of mind in which an individual is able to develop their potential, work productively, and creatively, and is able to cope with the normal stresses of life”,²¹ in students over 15 years in CSSNC stratified by gender identity. Furthermore, the study aims to compare coping strategies adopted to manage the health crisis by gender and to investigate associations between strategies adopted and a worsening of mental health due to the COVID-19 pandemic. It is essential to explore how girls and boys were coping during the pandemic and what coping strategies positively or negatively influenced changes in mental health to be able to understand the impacts of the pandemic and

to better inform appropriate school-based interventions, creating a safe and supportive environment that responds to their specific needs.

Method

Study design and participants selection

This is a cross-sectional study in 16 educational centres in Catalonia part of the CSSNC Project during 2021-2022.²⁰ Although it was an opportunistic sample, epidemiological, sociodemographic characteristics, and type of school (public, private, or chartered) were considered to assure heterogeneity. For the objectives of this study, students aged 15-19 years-old were included. The study was approved by the ethics committee of the IDIAP Jordi Gol i Gurina Foundation on 17 December 2020 (20/192-PCV). Informed consent was obtained from all the participants.

Data collection

Data collection took place from October-November 2021 through a questionnaire developed on the REDCap platform, or in paper when necessary. Each school was responsible to send a link containing the informed consent and the questionnaire available in Catalan, Spanish and English to teaching and non-teaching staff and students. Students over 15 years old signed the informed consent and completed the questionnaire by themselves. Response rate was 53% and 23% for staff and students, respectively.

Variables

The questionnaire including sociodemographic and health status variables was based on the WHO/Europe recommendation.²² The questions on impact of COVID-19 on psychological well-being of students and the coping strategies adopted were adapted from sections from the CASPE questionnaire.²³ The full list of variables is presented in [table 1](#).

Statistical analysis

Differences between proportions according to gender were compared using the Chi-Square, or the exact Fisher's exact test when more than 20% of cells have expected frequencies <5, and differences between means using the Student's t test. Only people identifying as men or women were included in the analysis because the small number of participants identified themselves as non-binary or other (12 people identified themselves as non-binary and 48 did not want to answer the question).

Logistic regression models were built to evaluate the association between coping strategies with self-perceived impact of the pandemic on mental health, adjusting by gender, house size and type of educational centre according to complexity. The dependent variable “self-perceived impact of the pandemic on mental health” was dichotomized as “it has improved or stayed the same” versus “it has worsened”. Adjusted OR (aOR) were calculated with a confidence interval of 95% (95%CI). The level of statistical significance was established as 0.05. All analyses were carried out using the R program, version 4.1.2.

Results

Of a total of 1171 participants, 626 (55.7%) identified as women, 478 (42.6%) identified as men, 12 (1.1%) as non-binary and 7 (0.6%) as other gender. Regarding complexity, 166 (14.2%) students were from low complexity schools, 140 (12.0%) medium low complexity,

Table 1

List of variables included in the questionnaire.

- *Sociodemographic variables*: type of educational centre (low, medium low, medium high, high or very high complexity), age, country of birth, level of studies (parents/tutors), change in household income since the start of the pandemic (increased, decreased, no change, unknown), living with people of 65 or over and/or with children and/or with healthcare professionals, house size. Gender identity options were female/male/non-binary/other. Female or male are also referred to as girl or boy below.
- *Health status*: self-perceived physical health and self-perceived mental health (excellent, very good, good, fair, poor), chronic illness, diagnosis for anxiety, and previous COVID-19 infection.
- *Self-perceived impact of the pandemic on mental health*: it has improved, it has worsened, and it has stayed the same.
- *Level of stress caused by uncertainty about the future generated by COVID-19*: Likert 7-point scale between 1 (less stressful) and 7 (more stressful).
- *Emotions felt in the last 7 days caused by the pandemic*: Likert 5-point scale between 1 (very slightly or not at all) and 5 (extremely). Emotions: anxiety/stress, fear, sadness, worry, irritability, loneliness, boredom, disappointment and frustration. An indicator was created with the sum of the answers from the 9 emotions lived (range 9–45) where higher values on the scale were related to worse psychological well-being.
- *Level of concern about the impact of the pandemic on different aspects of life in relation to the last 7 days*: Likert 5-point scale between 1 (very little or not at all) and 5 (a great deal).
- *Strategies used to cope with the pandemic*: participants had to select from a list coping strategies that were later classified as: 1) strategies related to social activities; 2) strategies related to solitary digital activities; 3) self-care strategies; 4) strategies related with unhealthy habits; 5) others, following previous studies using CASPE questionnaire.

Sources: WHO²² and Ladouceur²³.

700 (59.8%) medium high and the rest (14.1%) from high complexity schools.

The mean age of participating boys and girls was 16.9 and 16.8 years old, respectively, most were born in Spain (89.9% girls and 93.0% boys), and 9.3% of girls and 8.5% of boys reported a lower level of family income since the start of the pandemic. A higher proportion of boys compared to girls reported a university level of education for their mothers (36.4% boys and 29.7% girls) (Table 2).

Boys presented a better self-perceived general health compared to girls (95.3% good, very good or excellent for boys, and 92.3% for girls) and better self-perceived mental health (87.3% good, very good or excellent for boys, and 63.8% for girls). Additionally, the percentage diagnosed with anxiety before or after the pandemic started was higher in girls (20.6% girls and 7.3% boys) (Table 2).

A greater proportion of girls perceived a worsening in mental health due to the COVID-19 compared to boys (36.9% and 17.8%, respectively), while 5.7% of girls and 15.0% of boys perceived their mental health as improved in this same period ($p < 0.001$). Additionally, the level of stress caused by uncertainty about the future generated for COVID-19 was also greater in girls than in boys ($M = 4.1$, standard deviation [SD]: 1.7, and $M = 3.1$, SD: 1.8, respectively) (Table 3).

For all the emotions that can generate emotional discomfort, girls had statistically significant higher mean scores than boys, except for worry (Table 3). The main emotions reported for both girls and boys, respectively, were worry ($M = 2.7$, SD: 1.1, and $M = 2.2$, SD: 1.1) and boredom ($M = 2.7$, SD: 1.3, and $M = 2.4$, SD: 1.2), followed by frustration in girls ($M = 2.5$, SD: 1.3) and disappointment in boys ($M = 1.9$, SD: 1.1).

Most of the participants reported a high negative impact of the pandemic caused by the fact that friends or family could die (61.3% girls and 41.4% boys) and/or get sick (54.9% girls and 33.5% boys), together with the fear of losing jobs (themselves or family members) (46.5% girls and 29.6% boys) (Fig. 1).

One of the most frequent coping strategies used by adolescents to deal with the pandemic (Table 4) was contact with friends online and/or social media (76.5% and 68.6% in boys and girls, respectively), and other solitary digital activities like watching TV and/or films (80.8% and 69.7% in girls and boys, respectively). On the other hand, playing videogames was an activity mainly reported by boys in comparison with girls (78.7% and 28.1%, respectively). In addition, other social activities related to family such as reserving time to spend with people they lived with were more frequently reported by girls in comparison with boys (26.4% and 20.6%, respectively). More than half the adolescents also referred to self-care strategies like taking exercise (59.7% and 63.2% in girls and boys, respectively), and/or other individual non-digital activities such as reading and/or writing, higher among girls (41.2% girls and 14.8%

boys). Amongst unhealthy activities, girls reported smoking as a coping strategy more frequently than boys (9.9% girls and 5.4% boys), while eating less healthily (20.8% girls and 16.6% boys), and consumption of alcohol and/or drugs (11.2% boys and 8.1% girls) were reported similarly by gender. The percentage of girls who had spoken to a mental health professional was higher than boys (7.7% of girls and 2.6% of boys).

In the adjusted logistic regression analysis, we founded an association between the participation in family activities, such as playing games and/or sports, with a lower probability of declaring worsening of mental health due to the pandemic among girls (OR: 0.55; 95%CI: 0.38–0.79) (Table 5).

Playing board games, puzzles or crafting also showed a protective effect in relation to worsening of mental health among both girls and boys, respectively (OR: 0.62, 95%CI: 0.43–0.89 for girls and OR: 0.44, 95%CI: 0.22–0.88 for boys). Playing videogames also showed a protective effect in relation to worsening of mental health among girls (OR: 0.66, 95%CI: 0.45–0.97).

Among girls, other strategies related to unhealthy habits, like consumption of alcohol or drugs, were associated with a higher probability of declaring worsening of mental health due to the pandemic (OR: 2.61, 95%CI: 1.41–4.81). An association between smoking and a higher probability of declaring worsening of mental health due to the pandemic was seen among boys (OR: 2.45, 95%CI: 1.03–5.84).

Finally, adolescents who had spoken to mental health professionals reported a higher proportion of worsening of mental health (OR: 3.43, 95%CI: 1.82–6.46 for girls and OR: 4.56, 95%CI: 1.31–15.9 for boys) (Table 5).

Discussion

This study demonstrated the negative impact of the COVID-19 pandemic on psychological well-being in adolescents and a clearly worse impact on girls. The use of social and digital media to keep in contact with friends and reduce social isolation was the most commonly reported coping strategy for both girls and boys, whereas gender differences were seen in the use of other strategies used such as the higher proportion of girls using social activities related to family and/or friends in comparison to boys. This study also shows an association between coping strategies and self-perceived mental health status in adolescents in Catalonia. In this sense, the use of unhealthy habits during the early pandemic period (e.g., substance use) was associated with a higher probability of declaring worsening mental health among both girls and boys, whereas the use of other positive coping strategies (e.g., spending time with family) was associated with less adverse mental health among girls.

Table 2
Socio-demographic and health status characteristics of the sample according to gender identity.

Variables ^a	Girls (n = 626)		Boys (n = 478)		p
	n	%	n	%	
<i>Socio-demographics</i>					
Mean age (SD), years	16.9 (1.0)		16.8 (1.0)		0.102
Place of birth, Spain	551	89.9	436	93.0	0.076
<i>Level of education (fathers)</i>					
Primary or less	137	21.9	79	16.5	0.108
Secondary	235	37.5	179	37.4	
University	147	23.5	132	27.6	
Unknown or not applicable	107	17.1	88	18.4	
<i>Level of education (mothers)</i>					
Primary or less	106	16.9	47	9.8	0.002
Secondary	249	39.8	181	37.9	
University	186	29.7	174	36.4	
Unknown or not applicable	85	13.6	76	15.9	
<i>Level of education (guardians)</i>					
Primary or less	25	4.0	17	3.6	0.141
Secondary	69	11.0	46	9.6	
University	46	7.3	54	11.3	
Unknown or not applicable	486	77.6	361	75.5	
Size of home (m ²), more than 70 m ²	544	88.8	390	82.4	0.006
Living with children	296	48.4	236	51.0	0.412
Living with people > 65 years old	51	8.4	32	6.9	0.379
Living with healthcare professionals	96	15.7	59	12.7	0.167
Lower level of income (since COVID-19)	57	9.3	40	8.5	0.631
<i>Health status</i>					
Self-perceived health excellent/very good/good	550	92.3	444	95.3	0.045
Self-perceived mental health: excellent/very good/good	380	63.8	459	87.3	<0.001
Any chronic disease	60	10.1	40	8.6	0.411
<i>Diagnosis for anxiety</i>					
Never	473	79.4	432	92.7	<0.001
Yes, since the pandemic started	52	8.7	11	2.4	
Yes, before the pandemic	71	11.9	23	4.9	
Previous infection with COVID-19	128	20.4	82	17.2	0.151

SD: standard deviation.

^a Missing values not included in calculations (for all variables missing values were < 10%).**Table 3**
Level of stress caused by uncertainty about the future generated by COVID-19 and emotions felt by gender identity.


Variables	Girls (n = 626)		Boys (n = 478)		p
	Mean	SD	Mean	SD	
Self-perceived stress associated with the pandemic ^{a,b}	4.1	1.7	3.1	1.8	<0.001
<i>Emotions felt because of the pandemic^{a,c}</i>					
Anxiety or stress	2.4	1.3	1.7	1.0	<0.001
Fear	1.9	1.0	1.4	0.7	<0.001
Sadness	2.4	1.1	1.7	0.9	<0.001
Worry	2.7	1.1	2.2	1.1	0.255
Irritability	2.3	1.2	1.8	1.0	<0.001
Loneliness	2.1	1.2	1.6	1.0	<0.001
Boredom	2.7	1.3	2.4	1.2	0.044
Disappointment	2.3	1.3	1.9	1.1	<0.001
Frustration	2.5	1.3	1.8	1.0	<0.001
Total score	21.2	8.0	16.4	6.4	<0.001

SD: standard deviation.

^a Last 7 days.^b Likert scale from 1 (less stressful) to 7 (more stressful).^c Likert scale from 1 (very slightly or not at all) to 5 (extremely).

The higher percentage of girls than boys that perceived a worsening of mental health due to the pandemic and the higher level of stress caused by general uncertainty it produced is consistent with others studies.^{4–10} Emerging evidence indicates that girls perceive the impact of COVID-19 on their psychological well-being differently from boys.²⁴ However, a higher prevalence of diagnoses of anxiety since the start of the pandemic observed in the study (8.7% girls and 2.4% boys) suggests that objective data exists to confirm the worsening of mental well-being in girls. All adolescents experienced more intense emotions than usual linked to the pandemic such as: worry, boredom, frustration, and dis-

appointment, with girls presenting higher mean scores on most of the emotions that negatively affected psychological well-being. Other studies with children and adolescents in Spain during the first months of the pandemic revealed feelings like fear, sadness and boredom, typical of the break with routine and loss of predictability the pandemic supposed.⁷ Although no difference was observed in type of concerns most reported according to gender, there were differences in intensity with girls showing the highest prevalence, as reported previously.²⁵ The fact that family or friends could die if infected by SARS-CoV-2 was the commonest concern reported for both genders; a risk factor for poorer psychological



*p=0.05; **p<0.001

Figure 1. Level of concern (a lot/a great deal) about the impact of the pandemic on distinct aspects of life according to gender identity.

Table 4

Strategies used to cope with the pandemic by gender identity.

	Girls (n = 626)		Boys (n = 478)		p
	n	%	n	%	
<i>Social activities</i>					
Be in contact with friends online/social media	455	76.5	319	68.6	0.004
Participate in family activities	220	37.0	157	33.8	0.278
Reserve time to spend with people I live with	157	26.4	96	20.6	0.030
<i>Solitary digital activities</i>					
Watch TV/films	481	80.8	324	69.7	<0.001
Play videogames	167	28.1	366	78.7	<0.001
<i>Other non-digital solitary activities</i>					
Practice meditation/mindfulness/praying	69	11.6	19	4.1	<0.001
Writing/reading	245	41.2	69	14.8	<0.001
Play an instrument/listen to music	236	39.7	130	28.0	<0.001
Play board games, puzzles or crafting	195	32.8	102	21.9	<0.001
<i>Self-care strategies</i>					
Try to improve nocturnal rest	158	26.6	113	24.3	0.404
Do exercise	355	59.7	294	63.2	0.238
Eat more healthily	167	28.1	111	23.9	0.123
<i>Unhealthy activities</i>					
Eat less healthily	124	20.8	77	16.6	0.078
Drink alcohol and/or take drugs	48	8.1	52	11.2	0.085
Smoking	59	9.9	25	5.4	0.007
<i>Other strategies</i>					
Take vitamins/herbal remedies for immune system	46	7.7	16	3.4	0.003
Speak with mental health professionals	46	7.7	12	2.6	<0.001
Spend time in private outdoor areas	123	20.7	62	13.3	0.002
Follow a daily routine of activities	185	31.1	135	29.0	0.468

well-being in young people observed in several studies.²⁶ The second major stressor was fear of losing work, both for themselves and/or family members, consistent with a previous study in Catalonia where a decline in family income was associated with poorer overall well-being for girls and boys.²⁷

Amongst the most common strategies found in our study to cope with the pandemic, for both girls and boys, were maintaining contact with friends online or through social media, together with other solitary digital activities. Although that coping strategies alleviate the effects of social distancing, social network use has also been associated with negative psychological consequences in this context.²⁸

Girls reported in a higher proportion than boys the use of social activities related to family and/or friends as a coping strategy, as observed previously. In addition, among girls, spending time with family members was associated with less likelihood of declaring worsening mental health due to the pandemic.²⁹ Other studies also show that better communication and emotional support from family members during the pandemic was a protection against developing symptoms of anxiety or depression in children and adolescents.³⁰

As in other studies³¹ a large percentage of adolescents referred to self-care strategies like exercise to deal with the pandemic, which has shown mental health benefits for the young. However,

Table 5
Association between strategies to cope with the pandemic and worsening of mental health adjusted by gender identity.

	Girls (n = 626)			Boys (n = 478)		
	aOR	95%CI	p	aOR	95%CI	p
<i>Social activities</i>						
Be in contact with friends online/social media	1.18	0.79-1.76	0.430	1.03	0.61-1.74	0.923
Participate in family activities (games, sports, etc.)	0.55	0.38-0.79	0.001	0.69	0.41-1.17	0.171
Reserve time to spend with people I live with	0.70	0.47-1.03	0.07	0.78	0.41-1.46	0.436
<i>Solitary digital activities</i>						
Watch TV/films	0.75	0.55-1.31	0.467	0.85	0.51-1.42	0.538
Play videogames	0.66	0.45-0.97	0.036	1.32	0.70-2.49	0.387
<i>Other non-digital solitary activities</i>						
Practice meditation/mindfulness/praying	1.33	0.80-2.22	0.271	1.26	0.40-3.93	0.693
Writing/reading	1.23	0.87-1.73	0.242	0.75	0.36-1.55	0.441
Play an instrument/listen to music	1.31	0.93-1.86	0.122	1.25	0.74-2.10	0.402
Play board games, puzzles or crafting	0.62	0.43-0.89	0.011	0.44	0.22-0.88	0.021
<i>Self-care strategies</i>						
Try to improve nocturnal rest	1.08	0.74-1.58	0.676	0.96	0.54-1.70	0.895
Do exercise	1.33	0.94-1.88	0.107	0.74	0.45-1.20	0.223
Eat more healthily	1.01	0.69-1.47	0.964	0.90	0.51-1.61	0.731
<i>Unhealthy activities</i>						
Eat less healthily	1.26	0.84-1.91	0.268	1.20	0.65-2.24	0.557
Drink alcohol and/or take drugs	2.61	1.41-4.81	0.002	1.56	0.79-3.11	0.203
Smoke	1.70	0.98-2.94	0.058	2.45	1.03-5.84	0.043
<i>Other strategies</i>						
Take vitamins or herbal remedies for immune system	1.02	0.54-1.90	0.958	0.70	1.55-3.20	0.649
Speak with mental health professionals	3.43	1.82-6.46	<0.001	4.56	1.31-15.9	0.017
Spend time in private outdoor areas	0.90	0.59-1.37	0.611	0.74	0.33-1.63	0.451
Follow a daily routine of activities	0.86	0.59-1.24	0.419	1.03	0.61-1.75	0.911

aOR: odds ratio adjusted by house size and type of educational centre according to complexity; 95%CI: 95% confidence interval.

other less healthy habits like smoking, and consumption of alcohol and/or drugs, were also used as coping strategies.¹⁷ Compared to boys, more girls in our sample reported engaging in tobacco use to cope with the pandemic. This finding is consistent with evidence that girls may be more likely to use substances as a coping strategy and responds to their learning and development, as well as their mental health and psychosocial well-being needs.³² Worry and fear due to COVID-19, together with consumption to relieve feelings of emotional discomfort caused by anxiety, stress or depressive feelings, have been identified as factors associated with coping strategies involving substance use.³³

Finally, seeking professional help as a coping strategy was reported by a few number of adolescents, and more in girls, in line with previous studies showing young people will more often disclose their feelings to parents and friends than to professionals, this is in part due to mental health stigma and embarrassment, a lack of mental health knowledge and negative perceptions of help-seeking.³⁴

Some limitations of this study include a lack of pre-pandemic data to confirm a real decrease in mental health in adolescents using internationally validated scales. However, the questionnaire used was created specifically to measure perception of impact of the pandemic on psychological well-being in adolescents and was never intended to diagnose the real state of mental health of participants. On the other hand, as it is a transversal study, we cannot establish causality between worsening mental health and coping strategies adopted. Additionally, we cannot rule out a significant degree of memory bias, as in any retrospective survey. Finally, considering the non-probabilistic sampling method, the results cannot be generalized to the school population of Catalonia. However, one of the most important strengths of this study was the formation of a network of schools that worked together to develop, adapt and implement health protocols, studies and scientific dissemination in the face of the pandemic, providing safe attendance of face-to-face activities for students and education professionals. Still, in relation to the methodology, the study design, carried out concomitantly with the participatory research model, revealed a strengthening of science teaching and the strengthened student involvement, seek-

ing to mitigate as much as possible the impact of the pandemic on their mental health.

In conclusion, this study presents evidence on the negative impact of the COVID-19 pandemic on psychological well-being in adolescents in schools in Catalonia, especially in girls. Although the generalized use of social media technology as a coping strategy stands out, other activities and strategies have been employed, and differently by boys and girls. Some of these strategies have been related to better psychological well-being, but we must not forget that unhealthy strategies, such as consumption of alcohol and/or drugs and smoking, have also increased to deal with difficult situations. Due to the pandemic adolescents experienced much stronger negative emotions than usual such as worry, irritability or frustration; with girls presenting a higher proportion of them. More focus must be put on feelings generating emotional discomfort, especially if they are very intense and prolonged, because the absence of adequate tools to manage them can put those who experience them at risk. It is important to keep monitoring the medium- and long-term secondary impacts of the pandemic on mental health outcomes and consequent coping behaviours, especially considering gender perspective.

Finally, access for adolescents to emotional support services must be improved, both within schools and in specialized mental health centres, taking into account the need to support young people as they develop the necessary positive coping strategies required during health crises such as that of COVID-19.

Availability of databases and material for replication

All anonymized databases and their corresponding codebooks are available in the public repository <https://github.com/Escoles-Sentinella/Impact-on-psychological-wellbeing>. Due to legal restrictions in relation to the "Personal Information Protection Act", personal or spatial data that allow identified any participant, including the name of the school, which was used as an adjustment factor in the analysis, cannot be made publicly available. Requests for complementary data can be sent as a formal proposal to the CEEISCAT via email ceescat@iconcologia.net.

What is known about the topic?

The COVID-19 pandemic has affected the psychological well-being of adolescents who experienced high levels of stress and anxiety. In order to deal with the pandemic, adolescents used several coping strategies, such as physical activity and/or use of social networks to maintain contact with peers, among others.

What does this study add to the literature?

The study highlights gender differences in how the COVID-19 pandemic has affected adolescent mental wellbeing, and, for the first time in Spain, demonstrates a relationship between coping strategies used and self-perceived impact of COVID-19 on mental health by gender.

What are the implications of the results?

Findings highlights the need to monitor adolescent's mental health in schools considering a gender perspective. Support services for the development of healthy coping strategies during health crises like COVID-19 should be made.

Editor in charge

María del Mar García-Calvente.

Transparency declaration

The corresponding author on behalf of the other authors guarantee the accuracy, transparency and honesty of the data and information contained in the study, that no relevant information has been omitted and that all discrepancies between authors have been adequately resolved and described.

Authorship contributions

C. Folch, A. Colom-Cadena and J. Casabona made substantial contributions in the study conception and design. F. Ganem, I. Martínez and C. Cabezas made substantial contributions in data gathering, data analysis and interpretation. C. Folch wrote the first draft of the manuscript. All authors made substantial contributions in critical review with significant intellectual contributions. All authors approved the final version for publication.

Conflicts of interest

The authors would like to thank the Health Department and Department of Education of the Government of the Catalonia (Spain), the former Direcció General de Recerca i Innovació en Salut (DGRIS) and Institut Català de la Salut (ICS) that made this project possible, and all the health care professionals acting as a COVID-19 pandemic health taskforce in Catalonia. Also, we would like to specially thank the effort and dedication all the sentinel schools' staff, students and families who participated in the project.

Principal investigators: Jordi Casabona (Centre d'Estudis Epidemiològics sobre les Infeccions de Transmissió Sexual i Sida de Catalunya [CEEISCAT] – CIBERESP) and Josep Basora (Institut Universitari d'Investigació en Atenció Primària (IDIAP Jordi Gol)).

Project manager: Anna Bordas (CEEISCAT).

Technical committee: Jordi Casabona (CEEISCAT); Jordi Sunyer (ISGlobal); Antoni Soriano (Hospital Universitari Vall d'Hebron);

Rosina Malagrida (Living lab for Health, IRSICAixa) as Work Package coordinators. Cinta Folch, Pol Romano, Esteve Muntada, Anna Bordas, Fabiana Ganem, Andreu Colom i Jordi Casabona (CEEISCAT); Mireia Gascón, Maria Subirana, Pau Majo, Jordi Sunyer (ISGlobal); Rosina Malagrida, Laia Vives (Living lab for Health); Antoni Soriano, Pere Soler-Palacín (Hospital Universitari Vall d'Hebron).

Microbiology laboratories: Tomàs Pumarola, Andrés Antón, Cristina Andrés, Juliana Esperalba, Albert Blanco (Hospital Universitari Vall d'Hebron); Ignacio Blanco, Pere-Joan Cardona, María Victoria González, Gema Fernández, Cristina Esteban (Hospital Universitari Germans Trias i Pujol)

Data management and statistical analysis: Yesika Díaz, Lucia Alonso, Jordi Aceiton, Marcos Montoro (CEEISCAT).

Data protection officer and technical support: Esteve Muntada (CEEISCAT).

Communication manager: Pol Romano, Cristina Sánchez (CEEISCAT).

Field team: Maria Subirana, Pau Majo (ISGlobal); Jessica Fernández, Laia Vives (Living Lab for Health, IRSICAixa); Andreu Colom, Isabel Martínez, Marina Herrero, Alba García, Juan Rus, Paula Ribas, Alba Blanco (CEEISCAT).

Universitat Autònoma de Barcelona team: Juan Leyva, Mariela Patricia Aguayo, David Giménez i Carolina Watson.

Community paediatricians: Esperança Macià i Silvia Burgaya (CAP Manlleu); M. Teresa Riera-Bosch, Elisabet Sola (EAP Vic Nord); Lidia Aulet, María Mendoza, Lidia Busquets (EAP Vic Sud); Xavier Perramon, Júlia Sebastià (EAP Eixample Dret); Ana Moreno (Cap Ripollet); Xavier Duran, Belen Pérez (EAP Can Gibert del Pla); Anna Gatell (Equip Territorial de Pediatria Alt Penedès); Maria Coma (Hospital Universitari Joan XXIII).

Department of Health: Carmen Cabezas (Secretaria de Salut Pública de Catalunya, Departament de Salut); Ariadna Mas (Direcció Assistencial Atenció Primària, Institut Català de la Salut); Maria Antònia Llopis (Coordinació dels laboratoris de l'Institut Català de la Salut); Josep Vidal (Gerència Territorial de la Catalunya Central, Institut Català de la Salut); Sandra Pequeño and Jacobo Mendioroz (Subdirecció General de Vigilància i Resposta a Emergències de l'Agència de Salut Pública de Catalunya, Departament de Salut), Robert Fabregat (former Direcció General de Recerca i Innovació, Departament de Salut).

Department of Education: Josep González-Cambray (Conseller d'Educació); Núria Mora (Secretaria de Transformació Educativa); Maria Neus Fornells (former Gabinet del Conseller); Rut Ribas (former Direcció general de l'alumnat); Laia Asso (directora general d'Educació Inclusiva).

This work was supported by the Health Department of the Government of Catalonia with no grant number. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. F. Ganem, I. Martínez and A. Colom-Cadena received a salary from the abovementioned funder. None.

References

1. Nearchou F, Flinn C, Niland R, et al. Exploring the Impact of COVID-19 on mental health outcomes in children and adolescents: a systematic review. *Int J Environ Res Public Health*. 2020;17:8479.
2. Krishnaratne S, Pfadenhauer LM, Coenen M, et al. Measures implemented in the school setting to contain the COVID-19 pandemic: a rapid scoping review. *Cochrane Database Syst Rev*. 2020;CD013812.
3. Orgilés M, Morales A, Delvecchio E, et al. Immediate psychological effects of the COVID-19 quarantine in youth from Italy and Spain. *Front Psychol*. 2020;11:579038.
4. Wolf K, Schmitz J. Scoping review: longitudinal effects of the COVID-19 pandemic on child and adolescent mental health. *Eur Child Adolesc Psychiatry*. 2023;32:1–56.
5. Saule R, De Sario M, Bena A, et al. School closures and mental health, well-being and health behaviours among children and adolescents during the

- second COVID-19 wave: a systematic review of the literature. *Epidemiol Prev.* 2022;46:333–52.
6. Kauhanen L, Wan Mohd Yunus WMA, Lempinen L, et al. A systematic review of the mental health changes of children and young people before and during the COVID-19 pandemic. *Eur Child Adolesc Psychiatry.* 2023;32:995–1013.
 7. Amorós-Reche V, Belzunegui-Pastor A, Hurtado G, et al. Emotional problems in Spanish children and adolescents during the COVID-19 pandemic: a systematic review. *Clinica y Salud.* 2022;33:19–28.
 8. Theberath M, Bauer D, Chen W, et al. Effects of COVID-19 pandemic on mental health of children and adolescents: a systematic review of survey studies. *SAGE Open Med.* 2022;10, 20503121221086712.
 9. Kang S, Sun Y, Zhang X, et al. Is physical activity associated with mental health among Chinese adolescents during isolation in COVID-19 pandemic? *J Epidemiol Glob Health.* 2021;11:26–33.
 10. Halldorsdottir T, Thorisdottir IE, Meyers CCA, et al. Adolescent well-being amid the COVID-19 pandemic: are girls struggling more than boys? *JCPP Adv.* 2021;1:e12027.
 11. Magson NR, Freeman JYA, Rapee RM, et al. Risk and protective factors for prospective changes in adolescent mental health during the COVID-19 pandemic. *J Youth Adolescence.* 2021;50:44–57.
 12. Knight JA, Kehm RD, Schwartz L, et al. Prepubertal internalizing symptoms and timing of puberty onset in girls. *Am J Epidemiol.* 2021;190:431–8.
 13. Hoyt LT, Dotson MP, Suleiman AB, et al. Internalizing the COVID-19 pandemic: gendered differences in youth mental health. *Curr Opin Psychol.* 2023;52:101636.
 14. Folkman S, Lazarus RS. An analysis of coping in a middle-aged community sample. *J Health Soc Behav.* 1980;21:219–39.
 15. Golemis A, Voitsidis P, Parlapani E, et al. Young adults' coping strategies against loneliness during the COVID-19-related quarantine in Greece. *Health Promot Int.* 2022;37:daab053.
 16. Dvorsky MR, Breaux R, Cusick CN, et al. Coping with COVID-19: longitudinal impact of the pandemic on adjustment and links with coping for adolescents with and without ADHD. *Res Child. Adolesc Psychopathol.* 2022;50:605–19.
 17. Romano I, Patte KA, De Groh M, et al. Substance-related coping behaviours among youth during the early months of the COVID-19 pandemic. *Addict Behav Rep.* 2021;14:100392.
 18. Molero Jurado MDM, Pérez-Fuentes MDC, Fernández-Martínez E, et al. Coping strategies in the Spanish population: the role in consequences of COVID-19 on mental health. *Front Psychiatry.* 2021;12:606621.
 19. Marco-Ahulló A, Villarrasa-Sapiña I, Monfort-Torres G. Estudio descriptivo sobre las diferencias de género en el estrés académico derivado del contexto COVID-19 en población universitaria española. *Retos.* 2021;43:845–51.
 20. Bordas A, Soriano-Arandes A, Subirana M, et al. Study protocol for monitoring SARS-CoV-2 infection and its determinants in Catalonia (Spain): an observational and participatory research approach in a Sentinel Network of Schools. *BMJ Open.* 2022;12:e055649.
 21. Dhanabhakya M, Sarath M. Psychological wellbeing: a systematic literature review. *IJARST.* 2023;3:603–7.
 22. World Health Organization. Survey tool and guidance: rapid, simple, flexible behavioural insights on COVID-19. WHO. Regional Office for Europe; 2020. Available at: <https://apps.who.int/iris/handle/10665/333549>.
 23. Ladouceur C. Adolescent Symptom and Psychological Experience Questionnaire (CASPE). Pittsburgh, PA. 2020. Available at: <https://www.phenxtoolkit.org/toolkit/content/PDF/CASPE.Parent.pdf>.
 24. Pigaiani Y, Zocante L, Zocca A, et al. Adolescent lifestyle behaviors, coping strategies and subjective wellbeing during the COVID-19 pandemic: an online student survey. *Healthcare (Basel).* 2020;8:472.
 25. Hassan SN, Algahtani FD, Attaya MR, et al. The impact of extended e-learning on emotional well-being of students during the COVID-19 pandemic in Saudi Arabia. *Children.* 2021;9:13.
 26. Balluerka-Lasa N, Gómez-Benito J, Hidalgo-Montesinos M. Las consecuencias psicológicas de la Covid-19 y el confinamiento. Servicio de Publicaciones de la Universidad del País Vasco; 2020.
 27. Folch C, González-Casals H, Colom J, et al. Mental well-being during the COVID-19 confinement among adolescents in Catalonia: the role of demographic and other COVID-related variables. *Children.* 2022;9:783.
 28. Clark JL, Algoe SB, Green MC. Social network sites and well-being: the role of social connection. *Curr Dir Psychol Sci.* 2018;27:32–7.
 29. Riaz NA, Battista K, Duncan MJ, et al. Stronger together: coping behaviours and mental health changes of Canadian adolescents in early phases of the COVID-19 pandemic. *BMC Public Health.* 2023;23:319.
 30. Tang S, Xiang M, Cheung T, et al. Mental health and its correlates among children and adolescents during COVID-19 school closure: the importance of parent-child discussion. *J Affect Disord.* 2021;279:353–60.
 31. Ferguson KN, Coen SE, Tobin D, et al. The mental well-being and coping strategies of Canadian adolescents during the COVID-19 pandemic: a qualitative, cross-sectional study. *CMAJ Open.* 2021;9:E1013–20.
 32. Rogés J, Bosque-Prous M, Colom J, et al. Consumption of alcohol, cannabis, and tobacco in a cohort of adolescents before and during COVID-19 confinement. *Int J Environ Res Public Health.* 2021;18:7849.
 33. Rogers AH, Shepherd JM, Garey L, et al. Psychological factors associated with substance use initiation during the COVID-19 pandemic. *Psychiatr Res.* 2020;293:113407.
 34. Grove C, Marinucci A, Montagni I. Australian youth resilience and help-seeking during COVID-19: a cross-sectional study. *Behav Sci (Basel).* 2023;13:121.